



# VAJIRAM & RAVI

## Institute for IAS Examination

---

**UPSC CSE Prelims 2023**

## **Paper II – CSAT**

*for* Prelims 2023

The most detailed & authentic explanations to  
CSE Prelims 2023 - CSAT - II (Set C)

**UPDATED**

Answer Key  
&  
Detailed Explanations

# PowerUp

# Prelims Test Series

for **Prelims 2024**

- ✓ 30 GS Tests (15 Sectional, 5 Comprehensive, 10 Full Length)
- ✓ Detailed Model Answers & Explanation Videos
- ✓ Questions Based On UPSC Trend
- ✓ Equal Emphasis On Static & Current Events
- ✓ All India Ranking with AI-based Performance Analysis



**COMMENCES ON**  
**9th JULY, 2023**

## Prelims Assist - A Mentor Guided Prelims Test Series Programme



**30 GS Tests:**  
(15 Sectional,  
5 Comprehensive &  
10 Full-Length)



**Premier Mentor Sessions:**  
Learn straight from UPSC  
Rank Holders through  
exclusive monthly sessions



**One to One Mentor  
Support:** Get personalised  
guidance from subject  
Specific Mentors



**Targeted Feedback and  
Doubt Clearing:** Detailed  
Performance analysis &  
tailored feedback following  
each test



**Flexibility & Convenience:**  
Provision of opting Offline or  
Online Mentoring Sessions  
with Mentors



Avail access to  
exclusive content on  
the Student Portal  
(vajiramias.com)

**Starts 9 July 2023**

**Admission Open**



**COMMENCES ON**  
**9th JULY, 2023**

CSAT UPSC Prelims 2023 - Answer Key							
SET - C							
1. (b)	11. (a)	21. (d)	31. (c)	41. (d)	51. (a)	61. (b)	71. (d)
2. (b)	12. (d)	22. (c)	32. (c)	42. (a)	52. (a)	62. (a)	72. (a)
3. (b)	13. (c)	23. (d)	33. (b)	43. (b)	53. (a)	63. (d)	73. (c)
4. (a)	14. (d)	24. (c)	34. (a)	44. (c)	54. (c)	64. (d)	74. (b)
5. (c)	15. (b)	25. (a)	35. (d)	45. (d)	55. (c)	65. (d)	75. (a)
6. (b)	16. (b)	26. (c)	36. (d)	46. (c)	56. (c)	66. (a)	76. (d)
7. (a)	17. (a)	27. (d)	37. (d)	47. (a)	57. (a)	67. (d)	77. (c)
8. (c)	18. (a)	28. (d)	38. (a)	48. (a)	58. (c)	68. (a)	78. (a)
9. (b)	19. (d)	29. (c)	39. (c)	49. (a)	59. (b)	69. (a)	79. (d)
10. (c)	20. (c)	30. (d)	40. (a)	50. (b)	60. (b)	70. (c)	80. (b)

**Q1.**

**Answer: b**

**Q2.**

**Answer: b**

**Q3.**

**Answer: b**

**Q4.**

**Answer: a**

**Explanation:**

(i)  $7 \oplus 9 \oplus 10 = 8$

**Code :**  $7 + 9 + 10 = 26$  and  $2 + 6 = 8$

(ii)  $9 \oplus 11 \oplus 30 = 50$

**Code :**  $9 + 11 + 30 = 55$  and  $5 + 0 = 5$

(iii)  $11 \oplus 17 \oplus 21 = 13$

**Code :**  $11 + 17 + 21 = 49$  and  $4 + 9 = 13$

Therefore  $23 \oplus 4 \oplus 15$  will be coded as follows :

$23 + 4 + 15 = 42$  and  $4 + 2 = 6$ .

**Hence, option a.**

**Q5.**

**Answer: c**

**Explanation:**

For  $7x + 96$  to be divisible by  $x$ ,  $96$  should be divisible by  $x$ .

$96 = 2^5 \times 3$ , so it has 12 factors  $\{(5 + 1) \times (1 + 1)\}$

Therefore there are 12 possible values of  $x$ .

**Hence, option c.**

**Q6.**

**Answer: b**

**Explanation:**

Take the greatest possible four digits i.e 9, 8, 7 and 6 for p, q, r and s. (not in the same order)

$p + q + r + s = 30$  and  $(p + q)(r + s)$  is to be maximized the highest product of two numbers, whose sum is given, is obtained when both are equal.

Hence,  $15 \times 15 = 225$  is the greatest possible product.

{Like  $(9 + 6)(8 + 7)$ }

Hence, option b.

Q7.

Answer: a

Explanation:

1001 is a multiple of 13.

When a 3-digit number is multiplied with 1001, a six digit. Pattern, of the type 'abcabc' is formed.

So, if 999 is multiplied with 1001, it will make 999999, which is a multiple of 13.

If 99999999 ..... (99 times) is written, then every group of 6 nines will not give any remainder when divided by 13, as follows :

$$\underbrace{999999 | 999999 | 999999 | \dots}_{16 \text{ groups } (=96 \text{ digits})} \underbrace{999}_{\text{last 3 digits}}$$

Its only last three 9's which are responsible for any remainder.

$$\text{So, } \begin{array}{r} 13 \overline{) 999} \phantom{00} \\ \underline{-988} \phantom{00} \\ 11 \phantom{00} \end{array}; 11 \text{ is the remainder.}$$

Hence, option a.

Q8.

Answer: c

Explanation:

$$(11111111)^2 = 12345678987654321$$

Sum of its digits = 81

Hence, option c.

Q9.

Answer: b

Explanation:

Consider 90 two-digit numbers. (from 10 to 99)

$$\begin{aligned} \text{Sum of all the digits at units place} &= (1 + 2 + 3 + \dots + 9) \times 9. \\ &= 45 \times 9 = 405 \end{aligned}$$

$$\text{Sum of all the digits at tens place} = 10(1 + 2 + \dots + 9) = 450$$

Hundred's place = 1

$$\text{So, sum of all the digits} = 405 + 450 + 1 = 856$$


Hence, option b.

Q10.

Answer: c

Explanation:

No three of the six highlighted points are collinear.



$$\begin{aligned} \text{So, total number of triangles} &= {}^6C_3 \\ &= 20 \end{aligned}$$

Hence, option c.

**Q11.**

**Answer: a**

**Q12.**

**Answer: d**

**Q13.**

**Answer: c**

**Q14.**

**Answer: d**

**Explanation:**

Worst case analysis:

In this case Raj would be completing a red pair in the end. So, let's say Raj draws 18 white and 16 black shoes first and then 9 red shoes (all of left foot).

In this way Raj has drawn 44 shoes and a red pair is still not complete. The next shoe drawn will complete a red pair.

**Hence option d.**

**Q15.**

**Answer: b**

**Explanation:**

	6s	4s	1s
1	4	0	1
2	3	1	1
3	3	0	4

# ACE

Approach for  
Competitive Exams

## College Foundation Programme



140 G.S Class Lectures



40 Current Affairs  
Lectures



35 Lectures on Basic  
Numeracy, Logical  
Reasoning, Comprehension  
etc.



Printed study material  
& Monthly Current Affairs  
Magazine



Weekly Answer Writing  
Programme & MCQ  
Tests



Access to Revision  
Classes



Career Counselling &  
Guidance



Online/Offline Mentor  
Support Programme



**AVAILABLE IN CLASSROOM & LIVE-ONLINE MODE**

**COMMENCES ON  
30 JULY, 2023**

4	2	3	1
5	2	2	5
6	2	1	9
7	2	0	13
8	1	4	3
9	1	3	7
10	1	2	11
11	1	1	15
12	1	0	19
13	0	6	1
14	0	5	5
15	0	4	9
16	0	3	13
17	0	2	17
18	0	1	21
19	0	0	25

Therefore, 19 ways.

**Hence, option b.**

**Q16.**

**Answer: b**

**Explanation:**

- 1) Not possible. If 3 go in their respective envelopes, then fourth has to go in the correct envelope.
- 2) The number of ways of selecting (choosing) 2 letters out of 4 =  ${}^4C_2 = 6$  ways. These two letters will go in their correct envelopes. The remaining two can go in wrong envelopes only in one way (in each case).

Therefore, total 6 ways.

**Hence, option b.**

**Q17.**

**Answer: a**

**Explanation:**

$$100 = 2^2 \times 5^2$$

In the given number, there one are two multiplies of 5(85 and 95) and one multiple of 4 (96) which will make the product ending with double zeroes. Hence the remains will be zero.

Hence, option a.

**Q18.**

**Answer: a**

**Explanation:**

The given expression is  $(57242)^{945}$  945 is of the form  $4n + 1$ , therefore the units digit will be same as  $2'$ , which is 2.

**Hence, option a.**

**Q19.**

**Answer: d**

**Explanation:**

Consider the presentation

	A	B	C
+	D	E	F
1	1	1	1

Since, none of the six digits is zero,  $C + F$  must be equal to 11.

Now  $B + E$  must be equal to 10 {because 1 as carry-over from  $C + F$  will be added to make it 11}

Similarly  $A + D = 10$

Therefore  $A + D + B + E + C + F = 10 + 10 + 11 = 31$

Hence, option d.

Q20.

Answer: c

Explanation:

As per the given situation, the value of D would be 199. So  $\frac{199}{19} = 10\frac{9}{19}$  (least possible)

Therefore the difference between hundred's and units place = 8

Hence, option c.

Q21.

Answer: d

Q22.

Answer: c

Q23.

Answer: d

Q24.

Answer: c

Q25.

Answer: a

Q26.

Answer: c

Explanation:

1. Consider the (worst) case that the first 13 balls are black, next 19 are blue, next 25 are green, next 27 are yellow, next 37 are red, next 53 are white.

Till now  $180 - 6 = 174$  balls have been drawn and no complete group has been drawn. Now any ball drawn next will complete a group definitely. Hence this statement is correct.

2. Consider the case that first 54 balls are white, next 38 are red, next 28 are yellow, next 26 are green, next 20 are blue. Till now 166 balls have been drawn, yet all colours are not present.


The next ball drawn will fulfill the requirement. Hence this statement is correct.

Hence option c.

Q27.

Answer: d

Explanation:



Hence, option d.



Q28.

Answer: d

Explanation:

$C = D > F > A > B > E$

All five statements are needed.

Hence, option d

Q29.

Answer: c

Explanation:

Analysis of options:

(a)  $\boxed{D}$   
 $A = B; \downarrow$  {There's no connective established between B & E}  
 $\boxed{E} \leftrightarrow C$

Hence, not sufficient

(b)  $\boxed{C} \leftrightarrow \boxed{A} = \boxed{B}; \downarrow$  {There's no connective established between B & E}  
 $\boxed{E}$

Hence, not sufficient

(c)  $\downarrow$   
 $\boxed{E} \leftrightarrow \boxed{C} \leftrightarrow \boxed{A} = \boxed{B}$  {Using all form statements, it's evident that E is brother-in-law of B}

Hence, option c.

# Essay Course



Offers quality content & handouts that adhere to UPSC standards



Refines aspirants' writing ability



Evaluates proficiency with thorough tests




Focuses on various essential themes



COMMENCES ON  
11 JUNE, 2023

AVAILABLE IN CLASSROOM & LIVE-ONLINE MODE





**Q30.**

**Answer: d**

**Explanation:**

Option (D) is the answer as it contains one composite number (91), all the other options contain only prime numbers.

Hence, option d.

**Q31.**

**Answer: c**

**Q32.**

**Answer: c**

**Q33.**

**Answer: b**

**Q34.**

**Answer: a**

**Q35.**

**Answer: d**

**Q36.**

**Answer: d**

**Explanation:**

$$D(100A + 10B + C) = 3000 + 700 + 10D + D = 3700 + 11D$$

(i) If  $D = 4$ , the number  $[3700 + 11D]$  is divided by  $D (= 4)$

In this case, A, B and C are 9, 3 and 6 respectively.

(ii) If  $D = 5$ , the number  $[3700 + 11D]$  is divided by  $D (= 5)$

In this case, A, B and C are 7, 5 and 1.

Hence there's no unique value for  $(A + B + C)$

**Hence, option d.**

**Q37.**

**Answer: d**

**Explanation:**

Any number of the form 'XYZXYZ' is a multiple of 1001 (in a way that  $1001 \times XYZ = XYZXYZ$ )

So, irrespective of the values of X, Y and Z, the number XYZXZY is definitely divided by 1001.

$$1001 = 7 \times 11 \times 13.$$

**Hence option d.**

**Q38.**

**Answer: a**

**Explanation:**

The cubes in the question one found when all the surface cubes are removed (Let's call them core)


If the total number of smaller cubelets in  $n^3$ , then the number of core cubes is  $(n - 2)^3$ .

Here  $n = 5$ , so  $(n - 2)^3 = 27$ .

**Hence, option a.**

**Q39.**

**Answer: c**



**Q40.**

**Answer: a**

Let's say the total amount of work = 48 units.

Hence A would do 6 units a day, B would do 3 units a day and c would do 4 units a day.

Consider the following presentation:

Mon	Tue	Wed	Thu	Fri	Sat	Sun	Mon	Tue	Wed	Thu
A	B	C	A	B	C	A	B	C	A	B
6	3	4	6	3	4	6	3	4	6	3

48 units done in 11 days.

**Hence, option a.**

**Q41.**

**Answer: d**

**Q42.**

**Answer: a**

**Q43.**

**Answer: b**

**Q44.**

**Answer: c**

**Q45.**

**Answer: d**

Total marks deducted were 4.

Now, how these are going to be distributed among 4 papers are as follows.

Case 1 → When all 4 marks are deducted from 1 paper.

Total ways = 4

Case 2 → When 1 marks is deducted from each 4 paper.

Total ways = 1

Case 3 → When 3 marks is deducted from 1 paper & 1 marks is deducted from 1 paper.

Total ways = 12 (Since  $4p_1 \times 3p_1 = 12$ )

Case 4 → When 2 marks are deducted from any 2 papers.

Total ways = 6

Case 5 → When 2 marks are deducted from 1 paper & 1 marks are deducted from 2 papers.

Total ways = 12

So, overall ways =  $12+6+12+1+4=35$  ways.

**Hence, option d.**

**Q46.**

**Answer: c**

**Explanation:**

Four horizontal strips


← we can fill it in 3 ways

← again this can filled in 2 ways

← again in 2 ways

← again in 2 ways

So, depending upon question total number of ways in which we can fill the strips are


→  $3 \times 2 \times 2 \times 2 = 24$  ways

**Hence, option c.**

**Q47.**

**Answer: a**

**Explanation:**



So, maximum 6 files.

**Hence, option a.**

**Q48.**

**Answer: a**

**Explanation:**

Understanding from the question, there are two tasks - 1 and 2 and five persons - P, Q, R, S and T. As per the given constraints, the number of ways of assigning tasks to P and Q is just 1 (because P as well as Q can be given only Task -2)

No. of ways of assigning task to T is 2 (T can be given any of the two tasks)

No. of ways of assigning tasks to R and S is 3 ( R-1, S-2 ; R-2, S-1 ; R-2, S-2)

# Mains Test Series



Trains you to master 'The Art of Writing Answers'



Mentor Support & Detailed Feedback



Helps you to revise the syllabus comprehensively



Tests followed by Explanation Classes & Model Answer Papers



COMMENCES IN  
JUNE, 2023

**VTS EXCEL: 15+ Months**

**VTS UPGRADE: 9+ Months**

**VTS FOCUSED: 3+ Months**

**VTS SPRINT: 3+ Months**

**AVAILABLE IN  
OFFLINE & ONLINE  
MODE**

So, total number of ways of assigning tasks to five persons is  $1 \times 2 \times 3 = 6$

Hence, option a.

**Q49.**

**Answer: a**

**Explanation:**

According to question

1. Maximum number of coins required to buy

$$78 \text{ gm} = 7$$

↓

$$50 \text{ gm} \rightarrow 1 \text{ coin}$$

$$10 \text{ gm} \rightarrow 2 \text{ coins}$$

$$24 \text{ gm} \rightarrow 4 \text{ coins}$$

Statement 1 is correct.

2. To measure 78 gm, minimum we have to use 7 coins.

So, it is wrong.

Hence, option a.

**Q50.**

**Answer: b**

**Explanation:**

According to information given in question.

$$A + B \rightarrow A > B$$

$$A - B \rightarrow A \leq B$$

$$A \times B \rightarrow A \geq B$$

$$A \div B \rightarrow A < B$$

$$A \pm B \rightarrow A = B$$

Statement  $\rightarrow$  conclusion (we derived)

$$\left. \begin{array}{l} P \times Q \rightarrow P \geq Q \\ P - T \rightarrow P \leq T \\ T \div R \rightarrow T < R \\ R \pm S \rightarrow R = S \end{array} \right\} S = R > T \geq P \geq Q$$

**Conclusion:**

$$1. Q \pm T \rightarrow Q = T \rightarrow \text{wrong}$$

$$2. S + Q \rightarrow S > Q \rightarrow \text{correct}$$

Hence, option b.

**Q51.**

**Answer: a**

**Q52.**

**Answer: a**

**Q53.**

**Answer: a**

**Q54.**

**Answer: c**

**Explanation:**

Here principal and amount both are same over the period of same 1 year.

Since  $R$  (% annually) is compound half-yearly

and  $S$  (% annually) is compounded yearly

So, here amount is same, then

$S > R$

Hence, option c.

Q55.

Answer: d

Explanation:

Since 1186 is divided by the given no leaves remainder 31.

So,  $1186 - 31 = 1155 \leftarrow$  this will be exactly divisible.

$$\begin{array}{r} 5 \overline{)1155} \\ 11 \overline{)231} \\ 7 \overline{)21} \\ 3 \end{array}$$

Hence factors of 1155  $\rightarrow 5 \times 11 \times 7 \times 3$

Since, it is said that remainder is 31, so, numbers should be greater than 31, hence all those numbers are:

(1)  $3 \times 11 = 33$  (2)  $5 \times 7 = 35$  (3)  $5 \times 11 = 55$

(4)  $7 \times 11 = 77$  (5)  $15 \times 7 = 105$  (6)  $15 \times 11 = 165$

(7)  $15 \times 7 \times 11 = 1155$  (8)  $35 \times 11 = 385$  (9)  $77 \times 3 = 231$

Total such number are  $\rightarrow 9$ .

Hence, option d.

Q56.

Answer: c

Explanation:

Information given are:

$p < q < r$  &

$pp + qq + rr = \text{tto}$

it means we have choose all the values of  $p, q$  &  $r$  such that we get o, at unit digit of tto  $\leftarrow$  3 digit number.

Case-I : When  $p = 1$

	p	q	r		
(a)	1,	2,	7	$\rightarrow$	o
(b)	1,	3,	6	$\rightarrow$	o
(c)	1,	4,	5	$\rightarrow$	o

Case-II : When  $p = 2$

(a) 2, 3, 5  $\rightarrow$  o

Case-III : When  $p = 3$

(a) 3, 8, 9  $\rightarrow$  o

Case-IV : When  $p = 4$

(a) 4, 7, 9  $\rightarrow$  o

Case-V : When  $p = 5$

(a) 5, 6, 9  $\rightarrow$  o

1. Possible value of  $P = 5$  (1, 2, 3, 4 & 5)  $\rightarrow$  right

2. Possible value of  $Q = 6$  (2, 3, 4, 6 & 8)  $\rightarrow$  right

Hence, option c.

Q57.

Answer: a

Explanation:

All 4 digit numbers which are less than 2000 are  $\rightarrow$

1234

$$\begin{array}{r}
 1243 \\
 1324 \\
 1342 \\
 1423 \\
 + \quad 1432 \\
 \hline
 7998
 \end{array}$$

Hence, option a.

Q58.

Answer: c

Explanation:

Since, it is a case of circular arrangement but here  $n - 1$  will not be applicable.

So, number of selection will be in 12 ways.

Hence, option c.

Q59.

Answer: b

Explanation:

Since, it is based on calendar, so here we have to find out total number of odd days

So, divide  $10^{10}$  by 7, whatever, remainder we get that will be the number odd days.

$$\frac{10^{10}}{7} = \frac{(10^2)^5}{7} = \frac{(100)^5}{7} = \frac{2^5}{7} = \frac{32}{7} = 4 \text{ R } 4$$

Remainder is 4. So, if today is Sunday then on  $10^{10}$  day, it will be Thursday.

Hence, option b.

## Optional Enrichment Course



Mathematics



Sociology



Political  
Science & IR



Geography



Anthropology



History



Medical Science  
Crash Course



Value Addition  
Lectures



Targeted Study Material  
& Handouts



Course based on PYQs  
and Current Development



Answer Writing Practice  
with Detailed Discussions

COMMENCES IN  
OCTOBER, 2023

AVAILABLE IN  
CLASSROOM &  
LIVE-ONLINE MODE

**Q60.**

**Answer: b**

**Explanation:**

Duration of time after which each signal changes colours are:

$$1^{\text{st}} \rightarrow 25 \text{ sec}$$

$$2^{\text{nd}} \rightarrow 39 \text{ sec}$$

$$3^{\text{rd}} \rightarrow 60 \text{ sec}$$

Now, all three will become red after, take LCM of 25, 39, 60

$$5 \overline{)25, 39, 60}$$

$$5 \overline{)5, 39, 12}$$

$$5 \overline{)1, 39, 12}$$

$$5 \overline{)1, 39, 6}$$

$$5 \overline{)1, 39, 3}$$

$$5 \overline{)1, 13, 1}$$

$$1, 1, 1$$

$$\begin{aligned} \text{LCM} &= 5 \times 5 \times 2 \times 2 \times 3 \times 13 \\ &= 3900 \end{aligned}$$

So, all 3 will become red after 3900 sec.

Again, they will become green after 3900 sec.

So, total time from green to green

$$= 3900 + 3900 = 7800 \text{ sec}$$

$$= \frac{7800}{60} \text{ min}$$

$$= 130 \text{ min}$$

$$= 2 \text{ hr } 10 \text{ min}$$

So, they become again green at

$$2 \text{ pm} + 10 \text{ min}$$

i.e. 4:10 pm

**Hence, option b.**

**Q61.**

**Answer: b**

**Q62.**

**Answer: a**

**Q63.**

**Answer: d**

**Q64.**

**Answer: d**

**Explanation:**

Here we have to find remainder

$$\text{i.e. } \frac{2^{192}}{6} = \frac{2 \cdot 2^{191}}{6} = \frac{2^{191}}{3}$$

$$\Rightarrow \frac{(2)^{191}}{3} = \frac{(-1)^{191}}{3} = \frac{-1}{3} \equiv -2$$



$$\Rightarrow \frac{2}{3} \rightarrow \text{Remainder} = 2 \times \frac{2}{3} = 4$$

Because we have solve the fraction by 2.

Hence, option d.

**Q65.**

**Answer: d**

**Explanation:**

$$\begin{array}{cccc|cccc|cccc|cccc} A & B & C & \_ & \_ & A & B & C & \_ & D & A & B & B & C & D & \_ & A & B & C & D \\ \downarrow & \downarrow & & & & \downarrow & & & & & \downarrow & & & & & \downarrow & & & & \\ D & D & & & & C & & & & & A & & & & & & & & & \end{array}$$

Here, if we observe then in each slot two alphabet are repeated 2 are side by side.

Ex. D D | C C | B B | A A

Hence, option d.

**Q66.**

**Answer: a**

**Explanation:**

DEF

GHI

975

Here, E = 0, F = 8

D 0 8

So, 
$$\begin{array}{r} \text{GHI} \\ \hline 975 \end{array}$$

So, value of I = 7, H = 6

Now the value of D may be either 5 or 4

**Case-I :** when value of D is 4

then,  $xy = 408$

factors of  $408 = 2 \times 2 \times 2 \times 3 \times 17$

x and y are two digit numbers

so, pairs formed are

{ (i) 27 & 17, (ii) 34 & 12 }

$A + B + C = 1 + 2 + 3 = 6$

Since no digits are same

So, value of A = 1, B = 2, C = 3

**Case-II :** when value of D is 5

then,  $xy = 508$

factors of  $508 = 2 \times 2 \times \frac{127}{2}$

Since it is 3 digit

So, this will not be the case.

Hence, option a.

**Q67.**

**Answer: d**

**Explanation:**

Statement	Conclusion
True statement →	either P or Q get job
False statement →	neither R nor S get job
True statement →	maximum of two among R, S & T get the job
1 → wrong	
2 → wrong	
<b>Hence, option d.</b>	


**Q68.**

**Answer: a**

**Explanation:**

	Statement	Conclusion
I	→ If P → ✓, then Q & S → ✓	
II	→ If R & S → ✓, then T → ✗	

Now on the basis of above information



Now, when S is false then P is false.

∴ Either one or atleast P & R must be false.

So, it is true.

2. → It is wrong. As P is dependent on both Q & S.

**Hence, option a.**

# Prelims Exclusive General Studies Course



220 G.S Class Lectures &  
550+ Hours of Teaching



Complementary  
CSAT Classes



Printed Study Material  
called Yellow Books



Monthly Current Affairs  
Magazine



Access to Revision  
Classes



Online/Offline Mentor  
Support Programme



Regular Assessments  
and Objective Test Series



30 Prelims Test Series in  
G.S and Current Affairs




**AVAILABLE IN CLASSROOM & LIVE-ONLINE MODE**

**COMMENCES ON  
28 JULY, 2023**

Q69.

Answer: a

Explanation:



Total volume of cuboid =  $7 \times 5 \times 3 = 105 \text{ cm}^3$

Total number of cube of  $1 \text{ cm}^3 = 105$

1. No painted face

Total number of cube of  $1 \text{ cm}^3 = 5 \times 3 \times 1 = 15$

It is true.

2. Number of cubes of  $1 \text{ cm}^3$  with two faces painted (Blue & Green)

It will be 4

It is wrong.

Hence, option a.

Q70.

Answer: c

Explanation:

I N C O M P R E H E N S I B I L I T I E S  
T S S R P O N N M L I I H E E E C B

Only 2 positions are matching in both the cases.

Hence, option c.

Q71.

Answer: d

Q72.

Answer: a

Q73.

Answer: c

Q74.

Answer: b

Explanation:

The pattern is:

$1^{\text{st}}, 2^{\text{nd}}, 4^{\text{th}}, 7^{\text{th}}, 11^{\text{th}}, 16^{\text{th}}, \dots$   
 $\xrightarrow{+1} \xrightarrow{+2} \xrightarrow{+3} \xrightarrow{+4} \xrightarrow{+5} \dots$

Since the children are making a circle, the ring will come back to  $1^{\text{st}}$  child if the number in the pattern is  $(40x + 1)$ , like  $41^{\text{st}}, 81^{\text{st}}, 121^{\text{st}}$ , etc.

So,  $1 + (1 + 2 + 3 + \dots) = 40x + 1$ .

$\Rightarrow 1 + 2 + 3 + \dots + x = 40x$

$$\frac{x(x+1)}{2} = 40x$$

$x = 15$  does the work

Hence, option b.

Q75.

Answer: a

Explanation:

z z y y x x x x w w w w w ..... A.

In the above sequence, the total number of terms

$$= 2 + 3 + ..... + 27 = 13 \times 29 = 377$$

The middle term is 189<sup>th</sup> term.

Therefore  $2 + 3 + 4 + ..... x \leq 189$

$$x(x + 1) \leq 380$$

$x = 19$  fits the bill.

So, the 19<sup>th</sup> letter (starting from Z) would be 8<sup>th</sup> letter ( $27 - 19 = 8$ ) starting from A, which is H.

Hence, option a.

Q76.

Answer: d

Explanation:

St. 1.  $p \times q > 0$  establishes two things

(i) Neither of p or q is zero

(ii) Both have the same sign.

But doesn't tell us anything about values.

St. 2.  $p^2 > q^2$

Consider (p, q) as (4, 3) and (-4, -3). Both the pairs fit the bill but do not ascertain.

If  $p > q$  or  $q > p$ .

Even the combination of both the above statement doesn't decide if  $p > q$  or  $q > p$ .

**STEP UP**

**MENTORSHIP  
PROGRAMME**



Individual  
Mentorship



Prelims-focus  
Guidance



Mains-focused  
Guidance



Value Addition  
Lectures



COMMENCES ON  
12 JUNE, 2023

**AVAILABLE IN OFFLINE & ONLINE MODE**

Hence, option d.

Q77.

Answer: c

Explanation:

If  $p + q - r > p - q + r$

then  $q > r$ ; so p doesn't matter.

Therefore, neither statement 1 alone nor statement 2 alone will answer the question.

Combining both the statements:

$$\begin{array}{rcl} p + q > 0 & \Rightarrow & r > q \\ \hline r - p > 0 \\ \hline r - q > 0 \end{array}$$


Since  $q - r < 0$ , the answer is confirmed no.

Hence, option c.


Q78.

Answer: a

Explanation:



120 took T or C.



Total number of people =  $120 + x$

Milk only =  $x$

$\Rightarrow 120 + x = 5x$

$\Rightarrow x = 30$ ; sufficient

Hence, option a.

Q79.

Answer: d

Explanation:

St. 1: The digits are 1, 2, 3; but there would be 6 three-digit numbers possible with these digits.

Hence, option d.

Q80.

Answer: b

# Optional Test Series



Agriculture



Philosophy



Mathematics



Commerce  
& Accountancy



Public  
Administration



Political  
Science & IR



Economics



Psychology



Sociology



Geography



Anthropology



History



Medical Science



Trains you  
to master "The  
Art of Writing  
Answers"



Helps you to  
revise the syllabus  
comprehensively



Tests followed  
by Explanation  
Classes & Model  
Answer Papers

**COMMENCES IN JUNE, 2023**